Unit #1: Integration Review *Topic:* Area under the Curve *Objective: SWBAT solve various free response problems using their knowledge of integration.*

Warm Up #7:

1) Suppose $\int_{1}^{2} f(x)dx = -4$, $\int_{1}^{5} f(x)dx = 6$, $\int_{1}^{5} g(x)dx = 8$, and $\int_{5}^{9} g(x)dx = -2$, find a) $\int_{2}^{5} f(x)dx =$

b) $\int_{1}^{2} 3f(x) dx =$

c) $\int_{1}^{9} g(x) dx =$


2) For what value of k, k > 0 will the average value of $y = 4 - x^2$ on the closed interval [0,k] be -8?

OK, so let's put it all together now.

1) 2013 AB #4

The figure above shows the graph of f', the derivative of a twice-differentiable function f, on the closed interval $0 \le x \le 8$. The graph of f' has horizontal tangent lines at x = 1, x = 3, and x = 5. The areas of the regions between the graph of f' and the x-axis are labeled in the figure. The function f is defined for all real numbers and satisfies f(8) = 4.

- (a) Find all values of x on the open interval 0 < x < 8 for which the function f has a local minimum. Justify your answer.
- (b) Determine the absolute minimum value of f on the closed interval $0 \le x \le 8$. Justify your answer.
- (c) On what open intervals contained in 0 < x < 8 is the graph of f both concave down and increasing? Explain your reasoning.




2) 2012 AB #3


Let f be the continuous function defined on [-4, 3]whose graph, consisting of three line segments and a semicircle centered at the origin, is given above. Let g

be the function given by $g(x) = \int_{1}^{x} f(t) dt$.


- (a) Find the values of g(2) and g(-2).
- (b) For each of g'(−3) and g''(−3), find the value or state that it does not exist.



- (c) Find the x-coordinate of each point at which the graph of g has a horizontal tangent line. For each of these points, determine whether g has a relative minimum, relative maximum, or neither a minimum nor a maximum at the point. Justify your answers.
- (d) For -4 < x < 3, find all values of x for which the graph of g has a point of inflection. Explain your reasoning.



The function g is defined and differentiable on the closed interval [-7, 5] and satisfies g(0) = 5. The graph of y = g'(x), the derivative of g, consists of a semicircle and three line segments, as shown in the figure above.




(b) Find the x-coordinate of each point of inflection of the graph of y = g(x) on the interval -7 < x < 5. Explain your reasoning.

4) 2011 AB #4

The continuous function f is defined on the interval $-4 \le x \le 3$. The graph of f consists of two quarter circles and one line segment, as shown in the figure above.

Let $g(x) = 2x + \int_0^x f(t) dt$.

- (a) Find g(-3). Find g'(x) and evaluate g'(-3).
- (b) Determine the x-coordinate of the point at which g has an absolute maximum on the interval $-4 \le x \le 3$. Justify your answer.
- (c) Find all values of x on the interval -4 < x < 3 for which the graph of g has a point of inflection. Give a reason for your answer.



Graph of f


(d) Find the average rate of change of f on the interval $-4 \le x \le 3$. There is no point c, -4 < c < 3, for which f'(c) is equal to that average rate of change. Explain why this statement does not contradict the Mean Value Theorem.

5) 2014 AB #3

The function f is defined on the closed interval [-5, 4]. The graph of f consists of three line segments and is shown in the figure above.

Let g be the function defined by $g(x) = \int_{-3}^{x} f(t) dt$.

- (a) Find g(3).
- (b) On what open intervals contained in -5 < x < 4 is the graph of g both increasing and concave down? Give a reason for your answer.
- (c) The function h is defined by $h(x) = \frac{g(x)}{5x}$. Find h'(3).
- (d) The function p is defined by $p(x) = f(x^2 x)$. Find the slope of the line tangent to the graph of p at the point where x = -1.



Graph of f